
# **BEACH EROSION**









Nevis Historical and Conservation Society

Department of Physical Planning, Natural Resources and Environment, Fisheries Division University of Puerto Rico, Sea Grant College Program; Caribbean Development Bank UNESCO Environment and Development in Coastal Regions and in Small Islands

#### FORCES TO BE RECKONED WITH

**B**eaches are continuously changing – from day to day, month to month and year to year – as the natural forces of wind and water meet the land. These changes, which have been taking place for millions of years, are linked to variations in wind, waves, currents and sea level height.

**B**ut it is not just natural forces that change the beach, humans have a big role to play in this process as well, through mining stones and sand from the beaches and dunes, polluting and damaging coral reefs, and constructing buildings and walls too close to the sea.

Changes in the beaches affect everyone. The coast is a place we are all attracted to for recreation, sports and simple enjoyment. This constantly changing and hazard-prone coastal environment is also where the greatest financial investment is concentrated, as large tourism properties and establishments continue to be attracted towards the shores of St Kitts and Nevis. Tourism is a driving force in the country's economy so the state of its beaches is of major importance.


## Natural forces



- Hurricanes and tropical storms, occurring between June and November, cause dramatic beach changes usually resulting in serious beach erosion.
- High waves in winter resulting from storms in the North Atlantic Ocean, and known as swell waves, or locally as 'groundseas'.
- Sea-level rise, which is a long-term factor, taking place very slowly over decades causes shorelines to retreat inland.



**S**ince 1995, the Atlantic Basin (including the Atlantic Ocean, the Caribbean Sea, and the Gulf of Mexico) has entered a more active hurricane cycle, which may continue for more than 20 years.



Number of

Number of

Number of

Number of

Source: Gray et al http://typhoon.atmos.colostate.edu/forecasts/1999/nov99/

In the Atlantic Basin the number of really severe hurricanes (categories 3, 4 and 5) increased from one per year (1990 –1994) to four per year (1995 – 1999).

## Human forces

- Removing sand from beaches and dunes for construction purposes causes erosion and the loss of beaches and coastal lands, destroying the natural heritage of the coast and reducing the vibrancy of the tourism industry.
- Building too close to the beach interferes with the natural sand movement and may impede beach recovery after a serious storm or hurricane.
- Badly planned sea defences may cause the loss of the beach, and of neighbouring beaches.
- Pollution from human activities on the land may damage coral reefs and seagrass beds; these biological systems protect, and provide sand to the beaches.
- Removing vegetation from the dunes destabilises these
  protective sand barriers; and clearing sites inland results in
  increased soil and dirt particles being washed offshore and
  smothering coral reef systems.

Hurricane frequency between 1990 and 1999 in the Atlantic Basin

5 year periods

1990 – 1994 1995 – 1999


Pipes discharging onto the beach and into the sea cause pollution and may harm marine life, Pinney's Beach 2000




## WHAT'S HAPPENING WITH NEVIS' BEACHES?

In order to manage these changes, Nevis' beaches have been monitored since 1988 by the Nevis Historical and Conservation Society, who measure the beach slope and width every three months at numerous sites around the island.


Beach monitoring in progress at Hurricane Hill, 1988





## SAND IN, SAND OUT

**W**hen Hurricane Hugo struck in 1989, much of the beach was lost at Pinney's Beach on the west coast. In the months and years after the hurricane the beach recovered, but not to prehurricane levels. The beach was further eroded by successive hurricanes: Hurricane Luis in 1995, Hurricane Georges in 1998 and Hurricane Lenny in 1999.



BEACH PROFILE
Pinney's Cotton
Ground.
Beach erosion

Before Hurricane Hugo

(23/08/88)

After Hurricane Hugo

(17/10/89)

After Hurricane Luis

(15/10/95)



Pinney's Beach, August 1995, before Hurricane Luis



Pinney's Beach, October 1995, after Hurricane Luis


## **DUNES AS RESERVOIRS OF SAND**

Dunes function as reservoirs of sand, available for supply to the beach during storms. In addition, they protect coastal land from flooding.

BEACH PROFILE Indian Castle. Beach erosion

Before Hurricane Hugo (23/08/88)

After Hurricane Hugo (28/05/90)



At Indian Castle, there used to be extensive sand dunes, several metres high. These were mined extensively for construction sand in the 1970s and 1980s, and the remnant dunes were eroded by the recent hurricanes.


Extensive mining in the '70s and '80s destabilised the dunes at Indian Castle and left them vulnerable to hurricanes, 1988





## HERE TODAY, GONE TOMORROW

The table shows generalised rates of change at the measured beaches in Nevis. During the decade of the 1990s most of the beaches showed erosion; a result, at least in part, of the impact of several severe hurricanes since 1989. The erosion was especially severe on the west coast beaches.



Beach change rates in Nevis (metres per year)



A negative rate of change ( ) indicates erosion and retreat of the shoreline, a positive rate of change ( ) indicates accretion or advancement of the shoreline towards the sea.





Damaged beach bar at Golden Rock, Pinney's Beach, after Hurricane Luis, 1995

## WISE PRACTICES FOR A HEALTHY BEACH





This restaurant at Pinney's Beach was 30 metres from the sea when it was opened in August 1995. One month later it was destroyed by Hurricane Luis. The greater the distance from the sea, the safer the building

24m (80 ft) 37m (120 ft) 152m (500 ft) The state of the beach affects everyone's lives. There are no simple or universal solutions to shoreline erosion, since there are often several factors, both human and natural, contributing to the problem at a particular beach. Each beach behaves differently, so it is advisable to find out as much information as possible about a particular beach before taking any corrective action. It is necessary to consult the Department of Physical Planning, Natural Resources and Environment before undertaking any action at a beach.

Some forces of change, such as hurricanes and winter swells are natural, and there is little we can do to stop them, yet there are ways we can help to slow down the rate of erosion:

- Planning new development so that it is a 'safe' distance behind the beach will reduce the need for expensive sea defence measures in the future.
- Revegetating dunes with native vegetation e.g. grasses and vines, and planting beach areas beyond the reach of storm waves with salt-resistant, deep-rooting trees, such as seagrape.






- Resorting to 'hard' engineering structures such as seawalls, revetments and bulkheads, only when there is a need to protect beachfront property from wave action. Such structures, even with careful design, result in the loss or narrowing of the beach over time.
- Considering all other beach enhancement measures such as
  offshore breakwaters, groynes and beach nourishment (placing
  sand from the offshore zone or from an inland source on the
  beach) at a particular site. All such measures require careful
  design and environmental impact assessments, so always first
  consult the Department of Physical Planning, Natural Resources
  and Environment.



Planning officials inspecting a newly constructed rock revetment at Pinney's Cotton Ground, 1999


...but not in others (Nisbett 1992)

Groynes built along the north coast have stabilised the beach in some places (Newcastle 1997)...



By 2000, the beach had disappeared and land has been reclaimed and protected with a boulder revetment

## WISE PRACTICES / CHECKLIST



- Develop a proactive approach to coastal planning through the preparation of a coastal zone management plan.
- Involve all stakeholders (e.g. government agencies, coastal communities, nongovernmental organisations, coastal residents, beach users and others) in the preparation and implementation of the coastal zone management plan.
- Develop principles for coastal stewardship, so that everyone plays their role to the fullest.
- Maintain regular dialogue among all stakeholders, enhanced by awareness and education about beach and coastal issues.
- Review and carefully consider ALL options when planning ways to slow down the rate of coastline change, these should include planning, ecological and engineering measures.
- Continue to monitor the rate of coastline change and share the findings with all other stakeholders.
- Respect the rights of all beach users.
- Provide and clearly mark public accesses to all beaches.
- Conserve and restore vegetative cover, both adjacent to the beach in order to stabilise the sand, and further inland to reduce sediment reaching the reefs and seagrass beds.

For more information on shoreline change in **NEVIS** consult:

Nevis Historical and Conservation Society PO Box 563, Charlestown, Nevis, St Kitts & Nevis T: +1 869 469 5786/0408; F: +1 869 469 0274

Department of Physical Planning, Natural Resources and Environment Cotton House, Market Street, Charlestown Nevis, St Kitts & Nevis T: +1 869 469 5521; F: +1 869 469 5485 E: planevis@caribsurf.com For more information on shoreline change in the CARIBBEAN consult:

Coping with Beach Erosion by Gillian Cambers UNESCO Publishing, 1998 ISBN 93-3-103561-4

This booklet is a result of co-operation between UNESCO, the Caribbean Development Bank and Nevis' Governmental agencies

To view this booklet on line, please see: www.unesco.org/csi/act/cosalc/brochnev.htm